

Synchrony with Shunting Inhibition

Sachin S. Talathi^{1,*}, Dong-Uk Hwang¹, Abraham Miliotis¹, Paul R. Carney¹, and William L. Ditto¹

¹*J Crayton Pruitt Department of Biomedical Engineering,
University of Florida, Gainesville, FL 32611*

(Dated: January 30, 2009)

Spike time response curves (STRC's) are used to study the influence of synaptic stimuli on the firing times of a neuron oscillator without the assumption of weak coupling. They allow us to approximate the dynamics of synchronous state in networks of neurons through a discrete map, which can then be used to predict the stability of patterns of synchrony in the network. General theory for taking into account the contribution from higher order STRC terms, resulting from perturbations caused by synaptic stimuli lasting for more than one firing cycle of a neuron, in the approximation of the discrete map is still lacking. Here we present a general theory to account for higher order STRC corrections in the approximation of discrete map to determine the domain of 1:1 phase locked state in a network of two interacting neurons. We test the ability of this discrete map to predict the domain of 1:1 phase locked state in a network of two neurons interacting through a shunting synapse.

PACS numbers: 87.19.xm, 87.85.dm

Synchronous rhythms in the brain, in particular gamma rhythms (20-80 Hz) are known to constitute a fundamental mechanism for cognitive tasks such as object recognition[1, 2], associative learning[3, 4] and processing of sensory information in the brain [5, 6]. Early theoretical work by Wang and Rinzel [7], demonstrated the ability for reciprocally connected interneurons to exhibit synchronous rhythms, thereby providing a mechanistic framework for synchrony exhibited by a purely inhibitory neuronal network; i.e., neurons interacting through a hyperpolarizing GABA_A synapse ($E_R < V_{\text{rest}}$), where E_R is the reversal potential of GABA_A synapse and V_{rest} is the neuronal resting-potential. Recently there has been a resurgent interest in shunting inhibition ($V_{\text{rest}} < E_R \leq V_T$), where V_T is the threshold for a neuron to generate an action potential, following the experimental findings that GABA_A mediated inhibition in hippocampal slices of adult mammalian brains, is fast and shunting [8, 9]. Through simulation studies on a realistic network model of coupled interneurons, the authors in [10] have demonstrated that shunting enhances robust gamma-band synchrony in the network in the presence of moderately high heterogeneity. However due to the inherent complexity of the realistic network considered in [10], (a large network comprising of 200 interneurons, electrical coupling among neighboring pair of interneurons, the presence of synaptic propagation delay, fast synaptic decay time, strong synaptic conductance) it is difficult to elucidate the contribution of shunting to the enhancement of synchrony of the network in the gamma-band. Infact recent theoretical work [11] suggests that shunting may enhance stable asynchronous states in the network of coupled interneurons. However the authors

note that the observed difference in their analysis of synchrony induced through shunting synapse and those done by [10] may lie in the different regimes of synaptic coupling (weak versus strong) and the lack of heterogeneity in the intrinsic firing rates of the coupled neurons.

Here, we investigate whether the analytical framework of spike time response curves (STRC's) can be used to predict the synchronous state of 1:1 phase locking between two coupled neurons interacting through a strong shunting synapse in the presence of heterogeneity. We begin by demonstrating that synaptic stimuli through shunting inhibition to a neuron periodically firing in the gamma frequency band persists for 3 consecutive firing cycles as opposed to the case for a neuron receiving strong synaptic stimuli through a hyperpolarizing synapse. We then develop an analytical framework to approximate the dynamics of 1:1 synchronous state in a network of two neurons coupled via a strong shunting synapse through a discrete map by taking into account the higher order STRC contributions. We show that in the limit of zero higher order STRC contribution's the discrete map reduces to the well-known approximation for synchrony between two neurons coupled through a hyperpolarizing inhibitory synapse [12, 13, 14].

Each neuron in the network considered here is modeled based on a single compartment conductance based model for a fast spiking interneuron developed by [15]. The dynamical equation for the model neuron is given by

$$C \frac{dV(t)}{dt} = I_{DC} + I_S(t) + g_{Na} m_{\infty}^3 h(t) (E_{Na} - V(t)) + g_K n^4(t) (E_K - V(t)) + g_L (E_L - V(t)) \quad (1)$$

where $V(t)$ is the membrane potential, I_{DC} is the constant input DC-current that determines the intrinsic spiking period T_0 , E_r and g_r represent the reversal potential and conductance of ion channels with parameters

*Electronic address: sachin.talathi@bme.ufl.edu




Figure 1: (a) Schematic diagram demonstrating the effect of perturbation received by a spiking neuron at time t . The cycle containing the perturbation defines the first order STRC and the subsequent cycles define the higher order STRC terms. (b) The STRCs for hyperpolarizing synaptic input with $E_R = -80$ mV. (c) The STRCs for shunting synaptic input with $E_R = -55$ mV. The synaptic parameters are $\tau_R = 0.1$ ms, $\tau_D = 8$ ms, $g_s = 0.15$ mS/cm². The intrinsic period of ring for the neuron is $T_0 = 31$ ms and $V_{\text{rest}} = -65$ mV.

obtained from [15]. The inactivation variable $h(t)$ and the activation variable m_∞ for sodium channel and the activation variable $n(t)$ for the potassium channel satisfy the first order kinetic equations as described in [15]. $I_S(t) = g_s S(t)(E_R - V(t))$ is the synaptic current with E_R (mV) being the reversal potential of the synapse and g_s ($\frac{\text{mS}}{\text{cm}^2}$) being the strength of synaptic coupling. $S(t)$, gives the fraction of bound receptors and satisfy the following first order kinetic equation as described in [14, 16].

As a measure of the influence of synaptic stimuli on the firing times of a neuron, we define the spike time response curves (STRC's) $\Phi_j(t, \tau_R, \tau_D, g, E_R, T_0) = \frac{T_j - T_0}{T_0}$ [14, 17, 18], where T_0 is the intrinsic period of spiking, T_j represents the length of the j^{th} spiking cycle from the cycle $j = 1$ in which the neuron receives synaptic stimuli at time $0 < t < T_0$. The synaptic parameters are: τ_R : the synapse rise time, τ_D : the synaptic decay time, g : the synaptic strength and E_R : the reversal potential of the synapse. In general the synaptic input need not be weak. The STRC's are obtained numerically using the direct method [13] as shown in the schematic diagram in Figure 1a. The neuron firing regularly with period T_0 , is perturbed through an inhibitory synapse at time t after the neuron has fired a spike at reference time zero. The spiking time for neuron is considered to be the time when the membrane voltage V , crosses a threshold (set to 0 mV in all the calculations presented here). As a result of this perturbation, the neuron fires the next spike at time t_1 , representing the first cycle after perturbation of length T_1 , which is different from T_0 , the time at which




Figure 2: Schematic diagram to demonstrate the re-normalization and re-scaling procedure to determine the length of second cycle T_2 . Shown in red dashed lines are the effective spike times after the neuron receives two consecutive synaptic perturbations. Shown in black dotted lines is the change in the firing cycle caused by synaptic input in the first firing cycle. Shown in black dashed line is the unperturbed firing cycle for the neuron.




Figure 3: In (a) and (b) we show the color coded percent error $E = 100 \left| \frac{T_2^N - T_2^P}{T_2^N} \right|$ between the predicted value of T_2 , given by T_2^P through STRC's in equations 2 and 3 respectively and the actual value of T_2 given by T_2^N as determined by solving equation 1, numerically for neuron receiving two consecutive synaptic stimuli in successive firing. Inset shows the plot for the variation in T_2 (T_2^N shown in red and T_2^P shown in black) as a function of δt_2 for $\delta t_1 = 15$ ms

the neuron would have fired a spike in absence of any perturbation through inhibitory synapse. Depending on the properties of the synapse, i.e., synapse parameters: τ_R , τ_D and E_R ; through which the neuron receives the perturbing input, the length of subsequent cycles might change. For example, as can be seen from Figure 1b, for GABA_A mediated inhibition through synapse that is hyperpolarizing, the first order STRC is non-zero for all perturbation times $0 < \delta t < T_0$; the second order STRC is non-zero for $\delta t \rightarrow T_0$ and all higher order STRC

terms are zero. However for GABA_A mediated inhibition through synapse that is shunting (Figure 1c) the effect of perturbation lasts for the first three cycles including the perturbing cycle, i.e., both the first order and second order STRC's is non-zero for $0 < \delta t < T_0$ and the third order STRC is non-zero for $\delta t \rightarrow T_0$. Higher order STRC terms for shunting synapse are present because, shunting tends to depolarize the neuron thereby reducing the effective time for the occurrence of the next spike following synaptic stimuli. The asymptotic STRC Γ_∞ , gives the long term effect of perturbation received by a spiking neuron and is given through a linear sum of the first m non-zero STRC's, i.e., $\Gamma_\infty \stackrel{\text{def}}{=} \sum_{j=1}^m \Phi_j$. For hyperpolarizing synapse shown in Figure 1b, $m = 2$ and for shunting synapse shown in Figure 1c, $m = 3$. For all further calculations, unless otherwise mentioned, we will suppress the dependence of STRC on synaptic parameter's and the intrinsic period of the neuron T_0 . We further define the following two functions derived from STRC's: $R(\delta t) = T_0(1 + \Phi_1(\delta t)) - \delta t$ and $E_j(\delta t) = T_0(1 + \Phi_j(\delta t))$.

We will now use STRC's to develop a general framework to determine an approximate discrete map for 1:1 synchrony between two coupled interneurons interacting through a shunting synapse. We will begin by considering the simple case of a periodically firing neuron that receives synaptic stimuli through a shunting synapse in each of its two successive firing cycles at times δt_1 and δt_2 as shown in Figure 2. Our goal is to determine the length of the second cycle T_2 in this situation. In the presence of a single perturbation at δt_1 in the cycle 1, following from the definition of STRC's we have $T_2 = E_2(\delta t_1) = T_0(1 + \Phi_2(\delta t_1))$. Similarly in the presence of a single perturbation in cycle 2 at time δt_2 , again following from the definition of STRC's we have $T_2 = \delta t_2 + R(\delta t_2) = T_0(1 + \Phi_1(\delta t_2))$. In writing this equation we note that the default period of second cycle in the absence of the single perturbation at time δt_2 was T_0 . However if the synaptic input at δt_2 is followed by a synaptic perturbation in the previous cycle at δt_1 , the default length of the second cycle is no longer T_0 . Therefore, in order to correctly determine the length of second cycle in this case, we have to discount for the change in the default length of second cycle caused by synaptic input at time δt_1 . We do so by re-normalizing the synaptic perturbation time in the second cycle to $\delta t_2^e = \delta t_2 \frac{T_0}{E_2(\delta t_1)}$ and re-scaling the effective phase of perturbation by $\alpha_2^e = \frac{\delta t_2}{E_2(\delta t_1)} - \frac{\delta t_2}{T_0}$, as shown in Figure 2a. The re-normalization of the perturbation time and the re-scaling of the perturbing phase, allows us to discount for the effect of synaptic input at time δt_1 on the length of cycle 2. The length of cycle 2 now becomes $T_2 = \delta t_2 + R(\delta t_2^e) \cdot (1 - \alpha_2^e)$. In terms of STRC's we have,

$$T_2 = \delta t_2 + \left(T_0 \left(1 + \Phi_1 \left(\frac{\delta t_2}{1 + \Phi_2(\delta t_1)} \right) \right) - \frac{\delta t_2}{1 + \Phi_2(\delta t_1)} \right) \times \left(1 - \frac{\delta t_2}{T_0(1 + \Phi_2(\delta t_1))} + \frac{\delta t_2}{T_0} \right) \quad (2)$$

For a neuron receiving synaptic stimuli through a hyperpolarizing synapse, we have $\Phi_2(x) \approx 0$ (see Figure 1b). In this case, equation 1 reduces to

$$T_2 \approx T_0(1 + \Phi_1(\delta t_2) + \Phi_2(\delta t_1)) \quad (3)$$

The approximation in the form of equation 3, has been used in [19] to determine the effect of second order STRC component on stability of 1:1 synchronous state in a ring of pulse coupled oscillators; in [17] to determine phase resetting and phase locking in a hybrid circuit of one model neuron and one biological neuron and also recently in [18] to predict 1:1 and 2:2 synchrony in mutually coupled network of interneurons with synapse that is hyperpolarizing. In Figure 3a, we plot the percent error $E = 100 \left| \frac{T_2^N - T_2^P}{T_2^N} \right|$ between the predicted value for the length of second cycle: T_2^P , determined from equation 2, and the actual value: T_2^N , determined by numerically solving the ODE in equation 1, for a neuron receiving two consecutive synaptic stimuli at times δt_1 and δt_2 through a shunting synapse with following parameters: $\tau_R = 0.1$ ms, $\tau_D = 8$ ms, and $E_R = -55$ mV. In inset of Figure 3a, we plot T_2^N and T_2^P for the specific case of $\delta t_1 = 10$ ms. In Figure 3b, we show similar results from predicted value of T_2 estimated using equation 3. We see that while equation 2 is correctly able to predict the length of second cycle, equation 3 fails to capture the effect of second order STRC contribution in determining T_2 . We would like to emphasize that the expression for T_2 is only dependent on STRC's estimated for a given synapse type without any explicit assumption on the strength of synaptic input to the neuron and is valid both in the regime of weak and strong coupling and for slow and fast synaptic dynamics.

We can now generalize our approach of re-normalization and re-scaling to consider the situation when the neuron receives 3 synaptic stimuli in 3 consecutive firing cycles at time δt_1 in cycle 1, at time δt_2 in cycle 2 and at time δt_3 in cycle 3. This is an important case to consider in order to determine an approximate discrete map for 1:1 synchrony between two neurons interacting through a shunting synapse; since we know from Figure 1b that the effect of synaptic stimulus through a shunting synapse lasts for 3 consecutive firing cycles of the neuron. Following equation 2, the length of 3rd cycle in the presence of 3 consecutive synaptic stimuli can be written as: $T_3 = \delta t_3 + R(\delta t_3^e) \cdot (1 - \alpha_3^e)$; where $\delta t_3^e = \delta t_3 \frac{T_0}{E_3(\delta t_1, \delta t_2)}$ and $\alpha_3^e \approx \frac{\delta t_3}{E_3(\delta t_1, \delta t_2)} - \frac{\delta t_3 E_2(\delta t_1)}{(T_0)^2}$ where $\tilde{E}_3(\delta t_1, \delta t_2) = E_2(\delta t_2^e)$, represents the length of third cycle in the presence of two consecutive synaptic stimuli at times δt_1 and δt_2 . Note that although δt_3^e and α_3^e are dependent on timing of the occurrence of synaptic input in the first cycle, we have neglected the explicit contribution of $\Phi_3(\delta t_1)$ in the effective re-scaling term α_e^3 for determining T_3 . This is a reasonable approximation since $\Phi_3(\delta t) \approx 0$ for shunting synapse (see Figure 1b) resulting in $E_3(\delta t_1) \approx T_0$. Details on our estimate of the approximation for α_3^e is presented elsewhere [20].

We are now in the position to derive the approximate discrete map for 1:1 synchrony between neurons A and B firing with intrinsic period $T_0^A \neq T_0^B$ and coupled through a shunting synapse (see Figure 4a). The heterogeneity in the intrinsic firing rates of the two coupled neurons is quantified through $H = 100 \cdot \frac{I_{DC}^B - I_{DC}^A}{I_{DC}^A}$, where I_{DC}^A and I_{DC}^B are constant DC currents driving neurons A and B. From Figure 4b, when the two neurons are locked in stable 1:1 synchrony we have for neuron A, $t_{n+1}^A = t_n^A + \delta_n + R \left(\delta_n \frac{T_0^A}{\bar{E}_3(\delta_{n-2}, \delta_{n-1})} \right) \cdot (1 - \alpha_3^e)$, where t_n^X is the time of n^{th} for neuron $X = \{A, B\}$ and $\delta_n = \delta t_n^B - \delta t_n^A$. Since neuron B, does not receive any external perturbation, we have for neuron B, $t_{n+1}^B = t_n^B + T_0^B$. The discrete map for evolution of δ_n can then be obtained as:

$$\delta_{n+1} = T_0^B - R \left(\delta_n \frac{T_0^A}{\bar{E}_3(\delta_{n-2}, \delta_{n-1})} \right) \cdot (1 - \alpha_3^e) \quad (4)$$

The steady state solution to above equation can be obtained by solving for the fixed point δ^* of the discrete map defined by $\delta_{n+1} = \delta_n = \delta_{n-1} = \delta_{n-2} = \delta^*$. We then obtain $F(\delta^*) = T_0^B$, where $F(\delta^*)$ is given by

$$F(\delta^*) \approx \delta^* + R \left(\frac{\delta^* T_0^A}{\bar{E}_3(\delta^*, \delta^*)} \right) \times \left(1 - \frac{\delta^*}{\bar{E}_3(\delta^*, \delta^*)} + \frac{\delta^* E_2(\delta^*)}{(T_0^A)^2} \right) \quad (5)$$

In the limit of $\Phi_2 \approx 0$, $F(\delta^*) = T_0^A(1 + \Phi_1(\delta^*) + \Phi_2(\delta^*))$ corresponding to the well-known equation for the solution to the fixed point of discrete map for synchrony between two neurons coupled through a hyperpolarizing synapse [14]. Stability of the fixed point δ^* representing the solution to equation 5 requires $0 < \left| \frac{\partial F(\delta_n, \delta_{n-1}, \delta_{n-2})}{\partial(\delta_n, \delta_{n-1}, \delta_{n-2})} \right|_{\delta_n = \delta_{n-1} = \delta_{n-2} = \delta^*} < 2$. This stable fixed point then corresponds to the synchronous 1:1 locked state for the two coupled interneurons. In order to determine whether equation 4 can predict 1:1 phase locked states for the two neuron network interacting through a shunting synapse, we consider the specific case of neurons A and B coupled through a shunting synapse with parameters: $E_R = -55$ mV, $\tau_R = 0.1$ ms, and $\tau_D = 8$ ms. Neuron A receives fixed dc current I_{DC}^A , such that it is firing with intrinsic period of $T_0^A = 31$ ms. We solve equation 5, for different values of H , thereby modulating T_0^B , to determine the set of values for g_s , which will result in stable fixed point solution for equation 5. The solution is obtained by estimating STRC's for each value of g_s and then determining whether there is a fixed point solution to equation 5. In Figure 4c, we present the results of this calculation. For a given value of H , the curve in black gives the lower and upper bounds on the strength of coupling for shunting synapse g_s , for which a unique stable solution to equation 5 exists. For example with $H=50$, the range of values for g_s for which a unique stable solution exists




Figure 4: (a) Schematic diagram of the network considered. (b) Schematic diagram representing spike timing for neurons A and B when they are phase locked in 1:1 synchrony. In (c) and (d) we show domain of 1:1 synchrony estimated through STRC's from the discrete map in equation 5 (shown in black) and those obtained through numerical simulations of the network (shown in blue) for network of two interneurons coupled with shunting synapse with parameters $\tau_R = 0.1$ ms, $E_R = -55$ mV, $\tau_{uD} = 8$ ms, and $\tau_{uD} = 2$ ms respectively.

for equation 5 is $0.09 < g_s < 0.21$. This region of 1:1 synchronous locking is analogous to the classic Arnold tongue [14, 21], obtained for synchrony between two coupled nonlinear oscillators. Arnold tongue provides a two dimensional visualization of this dependence, as a bounded domain of region in the heterogeneity (H)-coupling strength (g) plane, where 1:1 synchrony between the two oscillators exist. In Figure 4b, the general feature of the Arnold tongue is represented as the region bounded by two black curves obtained through STRC by solving for fixed point of equation 5. In Figure 4b, shown in blue is a similar bound on the range of heterogeneity leading to synchronous oscillations between the two coupled neurons, obtained by numerically solving equation 1 for the evolution of the dynamics of the coupled neuron network. This curve is obtained by fixing the firing period of neuron A, T_0^A and varying the firing period of neuron B, by changing I_{DC}^B and determining the strength of synaptic coupling g_s that results in $\frac{T_0^B}{\langle T_0^A \rangle} \approx 1$. As can be seen from Figure 4b, the results match to those obtained through STRC calculations for fixed point of equation 5. In Figure 4d, we present similar calculation for the two neurons coupled through a fast shunting synapse with parameters: $E_R = -55$ mV, $\tau_D = 2$ ms, and $\tau_R = 0.1$ ms.

We note that the Arnold tongue showed in Figure 4c and Figure 4d are skewed to the right; i.e., $H > 0$, which suggests that shunting inhibition tends to promote 1:1 synchrony at higher frequencies of the driver neuron. Recently [14] have demonstrated the mechanism for phase locked state of 1:1 synchrony to exhibit identical synchrony, which is essential for the generation of synchronous oscillations in a larger network of neurons. It is therefore very likely then that through mechanism of

spike timing dependent plasticity [14], the two neurons may lock in identical synchrony at frequencies in the gamma range. This may result in the generation of gamma oscillations in larger network of interneurons interacting through shunting inhibition as has been demonstrated through simulation studies of realistic networks of neurons [10]. In conclusion, we have provided a general framework to study synchrony between neurons coupled through a shunting inhibition using discrete maps derived from spike time response curves.

-
- [1] T. Mima, T. Oluwatimilehin, T. Hiraoka, and M. Hallett, J Neurosci **21**, 3942 (2001).
 - [2] C. Baudry and O. Bertrand, TRENDS in Cog Sci **3**, 151 (1999).
 - [3] T. Gruber, A. Keil, and M. Muller, Neuroscience Letters **316**, 29 (2001).
 - [4] T. Gruber, M. Muller, and A. Keil, J Cog Neurosci **14**, 732 (2002).
 - [5] A. Engel and W. Singer, TRENDS in Cog Sci **5**, 16 (2001).
 - [6] F. Aoki, Clinical Neurophysiol **110**, 524 (1999).
 - [7] X. Wang and J. Rinzel, Neural Compute **4**, 84 (1992).
 - [8] M. Bartos, I. Vida, M. Frotscher, A. Meyer, H. Monyer, J. Geiger, and P. Jonas, Proc Natl Acad Sci **99**, 13222 (2002).
 - [9] I. Vida, M. Bartos, and P. Jonas, Neuron **49**, 107 (2006).
 - [10] M. Bartos, I. Vida, and P. Jonas, Nature Rev Neurosci **8**, 45 (2007).
 - [11] H. Jeong and B. Gutkin, Neural Compute **19**, 706 (2007).
 - [12] B. Ermentrout, Neural Compute **8**, 979 (1996).
 - [13] C. Acker, N. Kopell, and J. White, J Comp Neurosci **15**, 71 (2004).
 - [14] S. Talathi, D. Hwang, and W. Ditto, J Comp Neurosci **25**, 262 (2008).
 - [15] X. Wang and G. Buzsaki, J Neurosci **16**, 6402 (1996).
 - [16] H. Abarbanel, L. Gibb, R. Huerta, and M. Rabinovich, Biological Cybernetics **89**, 214 (2003).
 - [17] S. Oprisan, A. Prinz, and C. Canavier, Biophysical J **87**, 2283 (2004).
 - [18] S. Maran and C. Canavier, J Comp Neurosci **24**, 37 (2008).
 - [19] S. Oprisan and C. Canavier, Differential Equations and Dynamical Systems **9**, 243 (2001).
 - [20] S. Talathi, D. Hwang, and W. Ditto, J Comp Neurosci (Submitted).
 - [21] J. Kurths, A. Pikovsky, and M. Rosenblum, *Synchronization, a universal concept in non-linear science* (Cambridge University Press, 2001).